

SUCCESSFUL APPLICATIONS FIELD TRIALS


Canola—no fertilizer

Kelpgrow was applied on canola with different dilution rates:

1. Control (equivalent volume of water)
 2. Kelpgrow diluted 1:400 (2 Lt/ha)
 3. Kelpgrow diluted 1:200 (4 Lt/ha)
- and at various stages of growth: tillering, jointing, tillering and jointing.

No fertilizer was applied with seed at planting.

Results show a constant crop yield increase of the crop treated with Kelpgrow, ranging from +10% to +25%



Canola—with fertilizer


Kelpgrow was applied on canola with different dilution rates:

1. Control (equivalent volume of water)
 2. Kelpgrow diluted 1:400 (2 Lt/ha)
 3. Kelpgrow diluted 1:200 (4 Lt/ha)
- and at various stages of growth: tillering, jointing, tillering and jointing.

100 lbs of fertilizer (11-51-0) was applied with seed at planting.

Results show a constant crop yield increase of the crop treated with Kelpgrow, ranging from +26% to +32%

On Canola we suggest to apply our product at an early stage (late tillering—start of jointing) and using an application rate of 1 qt./acre.



West Coast Marine Bio Processing Corp.

#300 - 1497 Marine Drive, West Vancouver (BC), V7T 1B8 Canada

Tel. 604.338.8830, Fax. 604.608.5547, e-mail. info@marinebioprocessing.com

website: www.marinebioprocessing.com